



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

NOTE ON THE RADIAL VELOCITIES OF N. G. C. 3379 AND
N. G. C. 1700.

The object N. G. C. 3379 is of that class of small nebulae which photographs show to have an almost stellar condensation for a nucleus. The very faint trace of outlying nebulosity is insufficient for more than a guess concerning the spiral structure. A spectrogram of it was obtained in February, 1918, with the small focal-plane spectrograph attached to the 60-inch reflector. A total exposure time of 27^h45^m gave fair density and made measurements of radial velocity possible. Two measures by R. F. Sanford give +830 and +860 km/sec, respectively, when reduced to the Sun. Slipher has derived a value of +780 km/sec.

Another object of very similar appearance, N. G. C. 1700, has been photographed with the same spectrograph. The spectrogram is too much under-exposed to admit of trustworthy results, but a rough measurement reveals a high positive radial velocity. It will be of interest to see whether other objects of this type, of which several are known, will yield similar results.

FRANCIS G. PEASE.

A NINTH NOVA IN THE ANDROMEDA NEBULA

The photographs of the *Andromeda Nebula* made by Professor Ritchey on February 9th and February 10th, 1918, show a faint Nova in the following approximate position with respect to the nucleus:

$$\begin{aligned}\Delta\alpha &= +440'' \\ \Delta\delta &= +330''\end{aligned}$$

It is in an island of faint nebulosity in one of the dark rifts of the nebula.

The magnitude on February 9th was about 17.2 and on February 10th about 17.5.

The star does not appear on any other photograph of the nebula made at this observatory; the last preceding photograph was made January 15th, 1918, and the next following one was made July 6th, 1918. It is the ninth Nova to be found in this nebula. The eighth was discovered by Professor Ritchey on the first of the two plates on which the ninth appears.

JOHN C. DUNCAN.